

Association of T2 signal intensity of magnetic resonance imaging (MRI) of intracranial meningiomas with their consistency – a review

Thayane Araújo Lima¹, Davi Emanuel de Assis Fonseca¹, Luan de Sá Pinto Nóbrega Gadelha¹, Lucas Ravy Pereira Gomes de Souza¹, Keyvid dos Santos Pereira², Luiz Severo Bem Junior^{2,3}

¹Curso de Medicina da Faculdade de Ciências Médicas de Campina Grande – UNIFACISA, Campina Grande, Paraíba, Brasil.

²Departamento de Neurocirurgia do Hospital da Restauração, Recife, Pernambuco, Brasil.

³Programa de Pós-graduação de Neurociência, Universidade Federal de Pernambuco, Recife, Pernambuco, Brasil.

Abstract

Diferents studies have shown a possible association of neuroimaging as a predictor of intratumoral consistency, an important factor during surgery. To identify the correlation between the consistency of intracranial meningiomas and the image of the T2 sequence of magnetic resonance imaging. Using the PRISMA methodology, a search for clinical studies was carried out in the PubMed, Scielo, Medline and Cochrane databases; descriptors: “consistency”, “meningioma”, “MRI” and “prediction”. Twelve articles were found, with seven remaining after the inclusion criteria: articles written in English and published in the last ten years. The T2-weighted magnetic resonance sequence showed highest degree of correlation in the studies discussed, where T2 hyperintensity of soft tumors may be related to a higher water content, while T2 hypointensity for hard tumors may be due to the greater collagen and calcium content. Most studies allow an association to be established between the soft consistency of the tumor and signal hyperintensity in the T2 sequence of magnetic resonance imaging, with the consistency of the tumor being important, as the surgical difficulty and time depend on it. The hyperintensity of the lesion in T2 was associated with the soft consistency of the tumor, seen during the operation, whereas hypointense meningioma in T2 is associated with firm consistency.

Keywords: meningioma; consistency; MRI; prediction.

Resumo

Estudos diferentes têm demonstrado possível associação da neuroimagem como preditor da consistência intratumoral. Esta revisão tem o objetivo identificar a correlação entre a consistência dos meningiomas intracranianos e a imagem da sequência T2 da ressonância magnética. Utilizando a metodologia PRISMA, foi realizada busca por estudos clínicos nas bases de dados PubMed, Scielo, Medline e Cochrane; utilizando os descritores: “consistency”, “meningioma”, “MRI” e “prediction”. Doze artigos foram encontrados, restando sete após os critérios de inclusão: artigos escritos em inglês e publicados nos últimos dez anos. A sequência de RM ponderada em T2 mostrou o maior grau de correlação nos estudos, onde hiperintensidade em T2 dos tumores moles pode estar relacionada ao maior teor de água, enquanto a hipointensidade para os tumores duros pode ser devido ao teor de colágeno e cálcio. A maioria dos estudos permite estabelecer uma associação entre a consistência mole do tumor e hiperintensidade de sinal na sequência T2 da RM, sendo a consistência do tumor importante, pois ela é um fator influente na dificuldade e no tempo cirúrgico. A hiperintensidade da lesão em T2 foi associada à consistência mole do tumor, constatada no intra-operatório, já a hipointensidade em T2 associa-se a consistência firme.

Palavras-chave: meningioma; consistência; ressonância magnética; valor preditivo.

Introduction

For intracranial meningiomas, tumor consistency seems also to be of importance in determining the respectability and the surgical outcome, as well as in surgical planning¹, estimating the surgical time and the probability of using adjuvant therapy². Some studies claim that there is a probable association between the appearance of these tumors on magnetic resonance imaging (MRI) of the skull and intraoperative observations. They mainly used the T1 and T2 sequences to predict the operative findings and to compare the images of fluid attenuated inversion recovery (FLAIR) and the corresponding intraoperative tumor characteristics^{2,3,5,7,9}. However, previous studies vary widely in purpose, which refers to the central question of the article and quality, which refers to the method of work, the esteem and quality with which it was developed, and the accuracy of current neuroimaging techniques remains controversial with the theme. Controversies are about the negative correlation, denoted by some studies cited in the articles used as a reference^{3,4}

Therefore, the aim of the present study is to correlate the T2 signal intensity of magnetic resonance imaging (MRI) of intracranial meningiomas with its intraoperative consistency, in order to determine its predictive value to assist in planning and surgical outcome.

Methods

Study configuration

This study based its methods on the Main Items for Reporting Systematic Reviews and Meta-Analyzes (Prisma). Based on the question “Are there any association between T2 signal intensity in MRI and meningiomas consistency?”, that was established as a guiding principle, the articles selected by the search were submitted to an interpretative analysis.

Search strategy


On March 26, there was a search in pairs of the articles published in the following databases: PubMed, Scientific Electronic Library Online (SciELO), Medline and Cochrane, and later a complete text review was carried out by all researchers. The articles were selected from their year of publication, specifically from 2010 to 2020, and without the use of a secondary search from the references of the included articles.

The following keywords were used: meningioma AND consistency AND MRI AND prediction, in the form of the terms MeSH.

Selection criteria

For this systematic review, studies were selected based on the following criteria: articles written in English and with elapsed time of publication equal to or less than ten years, as reported in figure 1.

Figure 1. Description of the article selection process based on the eligibility criteria.



Using the selection criteria, as shown in the table above, studies published in English and in past 10 years, it was possible to select 9 articles that correlated the MRI sequences T1 and T2 to assess the correlation between the consistency of meningiomas in the preoperative period, consequently assisting the surgeon in preoperative planning and counseling. This correlation is established in each article referenced individually and explicitly throughout the text.

Results

Of the 9 selected studies, under the inclusion criteria: studies in English and published in the last ten years and exclusion criteria: greater than 10 years, which did not answer the question, 7 were case series. The case series totaled 737 patients. In this group, it was possible to perceive a positive and strong correlation regarding the ability to predict tumor consistency with sensitivity, mainly in the T2 sequence, in some studies with up to 90% sensitivity, and Smith *et al*³, who analysed one hundred cases in their series, demonstrated an accuracy 89% and positive predictive value of 84.9%, with $p < 0.05$, showing a useful and effective tool, on which the surgeon can base his decision for surgical planning.

Analyzing the studies included in this review, we will compare them to define whether there is a correlation between the consistency of the meningioma and the T2-weighted MRI sequence, as reported in chart 1.

Chart 1: Results of the research following the methods described in PRISMA.

Author and year	Methods	Patients analysed	Conclusion
Smith <i>et al</i> , 2016.	Case series	100 patients. Positive predictive value of 84,9%.	Propose a T2-based method of tumor consistency prediction with correlation to objective intraoperative consistency.
Watanabe <i>et al</i> , 2015.	Case series	43 patients. Sensitivity, specificity and accuracy were 89%, 79% and 81% for T2W; 89%, 76% and 79% for Flair; 100%, 74% and 79% for CE-FIESTA.	Suggest that a quantitative assessment using conventional T2W imaging or FLAIR may be a simple and useful method for predicting hard meningiomas.
Romani <i>et al</i> , 2014	Case series	110 patients.	Fractional Anisotropy (FA) value and Mean Diffusivity and FA maps are useful for prediction of meningioma consistency and, therefore, may be considered in the preoperative routine MRI examination of all patients with intracranial meningiomas.
Sitthina msuwan <i>et al</i> , 2012.	Case series	243 patients. T2W: $p=0.004$; FLAIR: $p=0.045$	Signal intensity on T2WI and FLAIR image can be used for insinuating meningioma

			consistency.
Alyaman <i>et al.</i> , 2018.	Case Series	70 patients. CUSA: p=0.046; FLAIR: 0.003	This study presents a new objective method to measure the consistency of intracranial meningiomas based on a simple algorithmic formula.
Hoover <i>et al.</i> , 2011.	Case Series	101 patients. Sensitivity: soft and firm were 90% and 56%, respectively (95% CI = 73-97% and 38-73%; p<0.001)	This tool using T1 and T2 series predicts meningioma consistency. Such knowledge should assist the surgeon in preoperative planning and counseling.
Karthigeyan <i>et al.</i> , 2019.	Case series	70 patients.	FLAIR hypointensity of meningiomas appears to have a significant independent association with the suboptimal operative plane with high specificity.

Smith *et al.*³, found from data collected from 100 patients, with tumors between 1.0-9.2 cm, 50 tumors were smooth, 29 intermediate and 21 were firm. The frequency of meningioma consistencies was divided into 3 groups: group A with tumor cerebellar/ peduncle index 72- weighted imaging intensity (TCTI)> 1.63; group B TCTI index between 1.33-1.63 and group C with TCTI index

<1.27. The comparison between the groups was done with 1-way ANOVA, with each TCTI index having statistical significance (p <0.00001). He states that the TCTI index correlates with the intraoperative consistency ratings. The author mentions that the consistency of meningioma is a continuum, with overlaps between the soft, intermediate and firm groups. However, the relevance is to differentiate the extreme groups (soft vs firm). Thus, using the ROC curve, they denoted a number for the definition of such consistencies, this optimal value being determined by a TCTI index of 1.41, with mild tumors greater than or equal to 1.41 and firm tumors less than 1.41. This value has a sensitivity of 81.9% and 84.9% specificity.

Chen *et al.*, mentioned in one of the analyzed articles⁴, who, in a prospective study of 54 patients, found a correlation of hyperintensity in T2 with mild tumors, but saw nothing of a correlation in T1; moreover, he says that multiple other studies have shown this same correlation.

According to Watanabe *et al.*⁵, from a study with 46 patients, in their analysis of correlation between the intensity signal index (SI) in the T2 image, FLAIR and CE-FIESTA showed a significant correlation, with p<0.05. On the other hand, no significant correlation was seen in T1 and CE-T1WI, with p> 0.05.

In the study by Romani *et al.*⁶, with 110 patients, it was shown that the hyperintense signal in T2WI is not predictive for the consistency of mild meningioma; in its case series, 40 firm and 16 mild tumors showed hyperintensity in T2WI.

Nevertheless, he states that, at the MD sequence, 39 firm tumors were isointense, and the hypointense signal correlates with statistical significance ($p = 0.008$) to firm tumors.

In the study selected for this review by Sitthinamsuwan *et al.*⁷, 243 patients were analyzed according to the consistency of the tumor and patient variables, such as age, sex, vasogenic edema and tumor characteristics in the T2 sequence of magnetic resonance, with the last variable reaching a 95% confidence interval (95% CI), indicating a statistic with a significant result of $p = 0.004$. With that, we evidenced, hypointense meningioma strongly related with hard consistency, while the tumor with higher signal intensity tended to be softer.

In the prospective observational study by Alyamany *et al.*¹, 54 patients were included in the total, as out of 70, 16 of them were excluded due to another diagnosis or incomplete data in order to measure the ratio of the ultrasonic cavitron surgical aspirator (CUSA) used in the resection of the tumors and the T2 sequence, with statistical significance of $p = 0.003$.

While Hoover *et al.*⁸, analyzed 101 patients regarding the use of both T1 and T2 in the preoperative period and showed that T2 hypodensity reached $p < 0.001$ in solid consistency and T2 hyperdensity $p = 0.05$ and the associated the isodensity or hypodensity of T1 this statistical significance becomes greater ($p < 0.01$), contributing to the objective of our research.

In this same perspective, Karthigeyan *et al.*⁹ showed that despite the extent of tumor vascularization, it was significantly correlated with the FLAIR intensity scores, in comparison with the T1 and T2 weighted images ($P < 0.05$) and that the relationship between image T2-meningioma-brain interface and the intraoperative plane was statistically significant ($P < 0.05$), being able to delimit the tumor well compared to brain tissue.

Discussion

Intracranial meningiomas account for 20–30% of all primary non-glioma intracranial tumors¹. Named by Harvey Cushing in 1922, it represents a tumor based on dural originating from arachnoid capillary cells². Epidemiologically, they occur in middle-aged adults and women are affected twice as often as men¹ with an incidence of approximately 2.3 per 100,000 for benign meningiomas and 0.17 per 100,000 for malignant meningiomas.

Meningiomas are well differentiated, benign and encapsulated lesions that press or that the brain measures that increases¹. They have a broad spectrum of histologically distinct clinical features and subgroups that are associated with a high risk of recurrence and morbidity, even after thorough inspection. The consistency of the tumor, its vascularization and the surgical plan are important factors².

The pillar of the treatment for meningiomas is the surgical resection that is usually achieved after devascularization the tumor from around the capsule then debunking it from inside using various tools including cavitron ultrasound aspirator

(CUSA) which is an instrument that combines suction and irrigation with ultrasound.

The correlation between the consistency of the meningioma and the imaging results, especially in MRI, is something that is much discussed today and with little consensus among the literature. Because of its surgical resolution, the consistency of the tumor is extremely important, as well as the size and location. It is already established that surgical difficulty is directly dependent on consistency, as well as increased surgical time in the case of firm consistencies. In addition, surgical time and the likelihood of adjuvant therapy can be estimated.

One of the studies found was a systematic review, in which n: 43 patients were evaluated, it was possible to corroborate the findings present in the other studies, showing that a predictive evaluation is useful to evaluate tumors. Finally, one of the studies, a literature review, went against the other studies, expressing the opinion that there is a need for more grounding in order to have this consistency direction with more reliability.

In the review by Yao *et al.*², 21 articles were included, of which 16 analyzed the usefulness of T1 and T2 to predict the consistency of meningioma. All concluded that T1 alone is not useful for such a prediction, and most noted that, if hyperintense in T2, the tumors tend to be mild. The probability is almost 100% that the tumor will be smooth, if it is hyperintense in T2 and hypointense in T1 consecutively. However, four authors cited by him did not find such a correlation with T2WI; 03 of

these observed only a parallel tendency of hyperintensity and hypointensity in T2WI for smooth and firm tumors, respectively, but without statistical significance.

Through a retrospective study of 54 patients, evidence was found of the relationship between meningioma consistency and T2 weighted image (T2WI MRI) findings, but, on the other hand, they did not do so following T1WI⁷. They also say that several other studies show this relationship between T2WI and consistency meningioma. In addition, it is worth mentioning that the hyperintensity on T2WI of soft tumors may be related to higher water content while the lower signal on T2WI for hard tumors might be due to less water and more collagen and calcium content. Based on the evidence above, despite some controversies and still lacking further studies, we found strong and concrete results that lead us to suggest the use of T2WI as a way to predict the consistency of meningioma⁷. This means that the hyperintensity in T2 means a high concentration of water and therefore appears shining in the image, resulting in a soft consistency tumor. Concomitantly, hypointensity in the same sequence shows the low amount of water, showing it to be a solid tumor.

However, in order to avoid researcher bias, we also mention studies that are opposed, that do not demonstrate this relationship and guide further studies on and also those articles that failed to establish a possible relationship between the variables studied, not validating the correlation between meningioma and T2WI³.

Finally, it is noteworthy that all studies of quantitative analysis of the review found a correlation of the T2WI sequence with the tumor consistency. The T2-weighted MRI sequence showed the highest degree of correlation in many studies discussed above as also talked about the correlation of consistency in the flair sequence in studies. Similarly, it is suggest and propose the use of both T2WI and the Flair sequence, to predict, in the preoperative period, the consistency of the tumor^{1, 2, 5, 7, 8}.

Conclusion

The consistency of meningioma has a predictive value for the surgical planning to be carried out and the respectability and the surgical outcome are determined¹. It is also possible to estimate the surgical time and the probability that the patient will need adjuvant therapy¹. The studies that were found are directed to very similar evidence regarding the relevance of the T2 sequence of MRI for determining the consistency of the meningioma in the preoperative period. The hyperintensity of the lesion in T2 was strongly associated with the soft consistency of the tumor¹, seen in the intraoperative period, while the T1 sequence has no predictive value². On the other hand, meningioma that is hypointense in T2 is associated with firm consistency⁸. From all that has been exposed, the association between the intensity of the meningioma signal in the T2 sequence and the consistency of the lesion is evident.

References

1. Alyamany M, Alshardan MM, Jamea AA, ElBakry N, Soualmi L, Orz Y (2018) Meningioma Consistency: Correlation Between Magnetic Resonance Imaging Characteristics, Operative Findings, and Histopathological Features. *Asian J Neurosurg*, 13(2):324-328.
2. Yao A, Pain M, Balchandani P, Shrivastava RK (2018) Can MRI predict meningioma consistency?: a correlation with tumor pathology and systematic review. *Neurosurg Rev*, 41(3):745-753.
3. Smith KA, Leever JD, Hylton PD, Camarata PJ, Chamoun RB (2017) Meningioma consistency prediction utilizing tumor to cerebellar peduncle intensity on T2-weighted magnetic resonance imaging sequences: TCTI ratio. *J Neurosurg*, 126(1):242-248.
4. Shiroishi MS, Cen SY, Tamrazi B, D'Amore F, Lerner A, King KS, *et al.* (2016) Predicting Meningioma Consistency on Preoperative Neuroimaging Studies. *Neurosurg Clin N Am*, 27(2):145-154.
5. Watanabe K, Kakeda S, Yamamoto J, Ide S, Ohnari N, Nishizawa S, *et al* (2016) Prediction of hard meningiomas: quantitative evaluate on based on the magnetic resonance signal intensity. *Acta Radiol*, 57(3):333-340.
6. Romani R, Tang WJ, Mao Y, Wang DJ, Tang HL, Zhu FP, *et al.* (2014) Diffusion tensor magnetic resonance imaging for predicting the consistency of intracranial meningiomas. *Acta Neurochir (Wien)*, 156(10):1837-1845.
7. Sitthinamsuwan B, Khampalikit I, Nunta-aree S, Srirabheebhat P, Witthiwej T, Nitising A (2012) Predictors of meningioma consistency: A study in 243 consecutive cases. *Acta Neurochir (Wien)*, 154(8):1383-1389.
8. Hoover JM, Morris JM, Meyer FB (2011) Use of preoperative magnetic resonance imaging T1 and T2 sequences to determine intraoperative meningioma consistency. *Surg Neurol Int*, 2:142.
9. Karthigeyan M, Dhandapani S, Salunke P, Singh P, Radotra BD, Gupta SK (2019) The Predictive Value of Conventional Magnetic Resonance Imaging Sequences on Operative Findings and Histopathology of Intracranial Meningiomas: A Prospective Study. *Neurol India*, 67(6):1439-1445.

Autor para correspondência
Thayane Araújo Lima
thayanearaujo1@gmail.com

Recebido: 26 de julho de 2020 Aceito: 11 de dezembro de 2020