



Review

Use of capsaicin as a model in the study of migraine: a literature review

Raisa Ferreira Costa¹  Emanuela Paz Rosas²  Daniella Araújo de Oliveira³  Marcelo Moraes Valença⁴ 

¹Postgraduate in Biological Sciences, Federal University of Pernambuco, Recife, Pernambuco, Brazil

²Keizo Asami Immunopathology Laboratory, Federal University of Pernambuco, Recife, Pernambuco, Brazil

³Department of Physiotherapy, Federal University of Pernambuco, Recife, Pernambuco, Brazil

⁴Department of Neuropsychiatry, Federal University of Pernambuco, Recife, Pernambuco, Brazil



Raisa Ferreira Costa
raissacosta@hotmail.com

Edited by

Juliana Ramos Andrade

Keywords:

Migraine
Pathophysiology
Capsaicin

Abstract

Capsaicin is able to induce mast cell degranulation, an event probably related to the pathophysiology of a migraine attack. The present review study aimed to address the mechanisms of action of capsaicin and other chemical inducers in mast cell degranulation and an interaction of nerves and events that happen in the dura mater with the activation of mast cells. A survey was carried out in the literature, from 1980 to 2019, in different databases, using the following terms: capsaicin, mast cell and dura mater. 36 articles were selected for this review. Studies indicate that the main mechanisms of action of capsaicin are chemical induction through the activation of TRPV1 channels, allowing calcium influx into neurons in the trigeminal ganglion of the dura mater, activating mast cell degranulation, releasing pro-inflammatory (e.g., histamine, oxide nitric) and vasoactive (e.g., CGRP and substance P) substances. Therefore, the use of capsaicin may be a tool to be used in an animal model to better understand the pathophysiology of migraine.

Received: January 14, 2021
Accepted: January 22, 2021

Introduction

The mast cells are present in the proximity of sensory nerve fibers in the rat dura mater.¹ There is an interaction between mast cells with C fibers in the dura mater.² Vasodilation, plasma leakage and degranulation of mast cells contribute to the release of pro-inflammatory substances in the dura mater, probably related to migraine attacks.³⁻⁵ The mast cell degranulation has been suggested as an important element of neurogenic inflammation, as mast cells exclusively release histamine when activated by the release of Calcitonin gene-related peptide (CGRP) at the trigeminal terminals.⁶⁻⁸ Histamine mediating CGRP receptors are likely to be located in mast cells.⁹⁻¹² In addition to CGRP, histamine also interacts with increased production of nitric oxide (NO) in the mast cells.¹³⁻¹⁶ All three chemical mediators contribute to vasodilation and neurogenic inflammation.¹⁷ Therefore, CGRP is able to release histamine and increase NO production from meningeal mast cells and act as a very potent vasodilator of intracranial arteries, occupying a prominent position in the nociceptive elements in the primary cascade of a headache attack.¹⁷

The capsaicin is one of the chemicals used in animal studies to activate the trigeminovascular system mimicking part of the migraine pathophysiological process and serving as an experimental model, as it is associated with the release of neuropeptides, through the opening of cationic channels that results in accumulation of intracellular calcium, and mast cell activation.¹⁸ These mast cells, through vasoactive mediators, contribute to stimulation of the large cranial arteries resulting in neurogenic inflammation and this is essential for their involvement in primary headaches.^{19,21} Thus, this review aims to deepen the knowledge about the mechanisms of action of capsaicin and other chemical inducers in the involvement of mast cell degranulation in a migraine attack.

Methods

This review was conducted with articles that used capsaicin to activate mast cells, in addition to other chemical inducers that demonstrated interaction between the dura mater and mast cells. The search was carried out with articles published between the years 1980 and 2019. Articles from SciELO, U.S. National Library of Medicine and the National Institutes Health (PubMed) and Web of Science databases were evaluated. As search strategy, the following terms were used: "Capsaicin" and "mast cell" and "dura mater" selected in consultation with the Medical Subject Headings (MeSH). The inclusion criteria were experimental model studies in rats that described the mechanisms of action of chemical inducers,

including capsaicin. Initially, the titles and abstracts of the articles found were read. Then, the articles were read and, finally, an additional search was carried out that tracked reference lists of all identified articles.

Results

Of the total 50 articles found, 36 were selected for this review.

Discussion

Capsaicin in experimental animal models is used to induce inflammation via chemical stress that increase vascular permeability in the dura mater by activating mast cells caused by the release of neuropeptides.^{22,23}

The transient potential of the cationic receptor of the V sub-family member 1 (TRPV1) also known as the capsaicin receptor, it's a channel that acts as a mediator to stress. After activation of these channels, they contribute to the opening of intracellular calcium and potassium output, leading to hyperpolarization of trigeminal neurons and vasodilation through the release of CGRP and substance P, neuromodulator that facilitates inflammatory processes. Thus, TRPV channels play vital roles in the vasculature and an important mediator for vascular responses.^{24,26} Baylie e Brayden²² demonstrated that stimulating actions increase the release of vasoactive peptides. Szolcsanyi²³ analyzed the depletion of sensory neurons by capsaicin and the relation of neuropeptides released in the analysis of the diameter of the cranial arteries, to verify whether there was dilation. Huang et al.²⁷ observe that chemical stimulation with capsaicin also influences the release of CGRP assessed immunohistochemically through the polyclonal antibody of rabbits.

The activation of mast cells by capsaicin leads to degranulation and the consequent release of pro-inflammatory substances (histamine and nitric oxide). Histamine increases CGRP levels released by C fiber neurons and consequently more numbers of degranulated mast cells, a positive feedback cycle, and nitric oxide contributes to controlling the viability of cranial arteries. In studies using capsaicin, it was observed that in neonates rats, chemical stimulation destroyed the peptidergic nerve endings, demonstrating that stresses in neonates do not induce mast cell degranulation.^{22,23}

The stimulating action of capsaicin to increase the release of neuropeptides causes depletion of sensory neurons, releasing

CGRP causing destruction of peptidergic nerve endings in neonatal animals that do not have a consistent neural structure. Dimtriadou et al. (1991) used capsaicin (50mg/kg; 48 h) administered subcutaneously in rats and do not had statistical difference in mast cell degranulation of the dura mater. Years later, with higher doses of capsaicin (50 mg/kg, 1 day of life and 100 mg/kg, 3 days of life) there was a statistical difference in the degranulation of mast cells of the dura mater, due to the increase in the capsaicin dose.^{28,29} Thus, neonates are not a good experimental model to mimic the pathophysiology of migraine, because they do not have a consistent and defined neural structure as in adult rats.

In studies with adult rats, capsaicin was able to induce a dose-response in the histamine release and consequent activation of mast cells, observing significant degranulation results.³⁰ As observed in another study that showed that foods containing capsaicin are able to suppress neuropeptides such as substance P that can cause the co-release of CGRP, since they are often located in the same neuron, thus activating the release of histamine in mast cells, mediating mast cell activation.³¹ This connection of mast cells with sensory neurons in the dura mater has a bidirectional response, where it signals responses with the nerves in relation to the external environment and endogenous and exogenous substances and degranulation in response to dural signals for the body's immunity.^{32,33}

In adult rats, it was also shown, through other inducers, as cholinergic agents (carbacol and nicotine); ovarian hormones and compound 48/80, a polymer that promotes histamine release, the percentage of degranulated mast cells in the dura mater and its relationship with migraine. It was observed that these chemical inducers had different mechanisms of action to activate mast cells, presenting significant results in the percentage of degranulation and in the pathophysiology of migraine.³⁴ In the study relating the ovarian hormones (estradiol and progesterone) to the density of mast cells in the dura mater and their potential role in headache, it is observed that the application of estrogen, but not progesterone, increases the density of mast cells.^{34,35} In the study of differentiating neuropeptides with vasoactive potentials, PACAP-38 to PACAP-27, in the dura mater and the frequency of mast cell degranulation, it was found that the difference is not in the pituitary adenylate cyclase-activating polypeptide type I (PAC1) receptor but because PACAP38 has greater induction in mast cell degranulation by acting via phospholipase C, which presents enzymes that contribute to the opening of intracellular calcium.³⁵ And in the study of cholinergic agents in the activation of calcium channels sensitized by CGRP, there is a relationship that suggests nociceptive regulation of the meninges, providing firing to primary afferents assuming

that this is the source of the headache, causing a significant mast cell degranulation and inducing vasodilation in the dura mater.³⁶ Then, other stimulants have mechanisms of action that resemble capsaicin to degranulate mast cells and increase the levels of these three inflammatory mediators.

Thus, probably due to the actions of capsaicin, it is possible to study the pathophysiology of the migraine by mast cell degranulation. Mast cell degranulation was considered a fundamental element for migraine physiopathology, as activation and consequent degranulation increases the plasma levels of three chemical mediators (CGRP, histamine and nitric oxide) that are related to vasodilation. Capsaicin, with its neuron-stimulating action, opens calcium ion channels, leading to neuron hyperpolarization and the release of vasoactive neuropeptides, activating mast cells and causing vasodilation.

Conclusion

The method of inducing mast cell degranulation by chemical activation of capsaicin, and other inducers, mimics the pathophysiological process of migraine by activating the trigeminovascular system and neurogenic inflammation.

Raisa Ferreira Costa

<https://orcid.org/0000-0002-1180-6689>

Emanuela Paz Rosas

<https://orcid.org/0000-0001-9895-5654>

Daniella Araújo de Oliveira

<http://orcid.org/0000-0002-6013-978X>

Marcelo Moraes Valença

<https://orcid.org/0000-0003-0678-3782>

References

1. G. Barbara, V. Stanghellini, R. De Giorgio, C. Cremon, G. S. Cottrell, D. Santini, G. Pasquinelli, A. M. Morselli-Labate, E. F. Grady, N. W. Bunnett, S. M. Collins and R. Corinaldesi. **Activated mast cells in proximity to colonic nerves correlate with abdominal pain in irritable bowel syndrome.** *Gastroenterology* 2004;126(3):693-702 Doi: <https://www.doi.org/10.1053/j.gastro.2003.11.055>
2. F. L. Rice, J. Y. Xie, P. J. Albrecht, E. Acker, J. Bourgeois, E. Navratilova, D. W. Dodick and F. Porreca. **Anatomy and immunochemical characterization of the non-arterial peptidergic diffuse dural innervation of the rat and Rhesus monkey: Implications for functional regulation and treatment in migraine.** *Cephalalgia* 2017;37(14):1350-1372 Doi: <https://www.doi.org/10.1177/0333102416677051>
3. P. Geppetti, E. Rossi, A. Chiarugi and S. Benemei. **Antidromic vasodilatation and the migraine mechanism.** *J Headache Pain*

- 2012;13(2):103-111 Doi: <https://www.doi.org/10.1007/s10194-011-0408-3>
4. H. E. Boran and H. Bolay. **Pathophysiology of Migraine.** *Noro Psikiyatr Ars* 2013;50(Suppl 1):S1-s7 Doi: <https://www.doi.org/10.4274/Npa.y7251>
 5. S. E. Erdener and T. Dalkara. **Modelling headache and migraine and its pharmacological manipulation.** *Br J Pharmacol* 2014;171(20):4575-4594 Doi: <https://www.doi.org/10.1111/bph.12651>
 6. H. Dong, X. Zhang and Y. Qian. **Mast cells and neuroinflammation.** *Med Sci Monit Basic Res* 2014;20(1):200-206 Doi: <https://www.doi.org/10.12659/msmbr.893093>
 7. A. C. Rosa and R. Fantozzi. **The role of histamine in neurogenic inflammation.** *Br J Pharmacol* 2013;170(1):38-45 Doi: <https://www.doi.org/10.1111/bph.12266>
 8. K. B. Alstadhaug. **Histamine in migraine and brain.** *Headache* 2014;54(2):246-259 Doi: <https://www.doi.org/10.1111/head.12293>
 9. T. C. Theoharides, J. Donelan, K. Kandere-Grzybowska and A. Konstantinidou. **The role of mast cells in migraine pathophysiology.** *Brain Res Brain Res Rev* 2005;49(1):65-76 Doi: <https://www.doi.org/10.1016/j.brainresrev.2004.11.006>
 10. A. F. Russo and I. M. Dickerson. **CGRP: a Multifunctional Neuropeptide.** In: A. Lajtha and R. Lim, editors. *Handbook of Neurochemistry and Molecular Neurobiology: Neuroactive Proteins and Peptides.* Boston, MA: Springer US; 2006. p. 391-426.
 11. N. Schwenger, M. Dux, R. de Col, R. Carr and K. Messlinger. **Interaction of calcitonin gene-related peptide, nitric oxide and histamine release in neurogenic blood flow and afferent activation in the rat cranial dura mater.** *Cephalalgia* 2007;27(6):481-491 Doi: <https://www.doi.org/10.1111/j.1468-2982.2007.01321.x>
 12. J. K. Lennerz, V. Rühle, E. P. Ceppia, W. L. Neuhuber, N. W. Bunnett, E. F. Grady and K. Messlinger. **Calcitonin receptor-like receptor (CLR), receptor activity-modifying protein 1 (RAMP1), and calcitonin gene-related peptide (CGRP) immunoreactivity in the rat trigeminovascular system: differences between peripheral and central CGRP receptor distribution.** *J Comp Neurol* 2008;507(3):1277-1299 Doi: <https://www.doi.org/10.1002/cne.21607>
 13. L. Edvinsson. **Sensory nerves in man and their role in primary headaches.** *Cephalalgia* 2001;21(7):761-764 Doi: <https://www.doi.org/10.1177/033310240102100705>
 14. N. Suzuki, Y. Fukuuchi, A. Koto, Y. Naganuma, K. Isozumi, S. Konno, J. Gotoh and T. Shimizu. **Distribution and origins of cerebrovascular NADPH-diaphorase-containing nerve fibers in the rat.** *Journal of the Autonomic Nervous System* 1994;49(Suppl): 51-54 Doi: [https://doi.org/10.1016/0165-1838\(94\)90086-8](https://doi.org/10.1016/0165-1838(94)90086-8)
 15. F. Tore, O. T. Korkmaz, D. Dogrukol-Ak and N. Tunçel. **The effects of vasoactive intestinal peptide on dura mater nitric oxide levels and vessel-contraction responses in sympathectomized rats.** *J Mol Neurosci* 2010;41(2):288-293 Doi: <https://www.doi.org/10.1007/s12031-009-9310-8>
 16. E. Kilinc, T. Firat, F. Tore, A. Kiyani, A. Kukner and N. Tunçel. **Vasoactive Intestinal peptide modulates c-Fos activity in the trigeminal nucleus and dura mater mast cells in sympathectomized rats.** *J Neurosci Res* 2015;93(4):644-650 Doi: <https://www.doi.org/10.1002/jnr.23523>
 17. D. Koyuncu Irmak, E. Kilinc and F. Tore. **Shared Fate of Meningeal Mast Cells and Sensory Neurons in Migraine.** *Front Cell Neurosci* 2019;13(1):136 Doi: <https://www.doi.org/10.3389/fncel.2019.00136>
 18. J. Donnerer and R. Amann. **Capsaicin-evoked neuropeptide release is not dependent on membrane potential changes.** *Neurosci Lett* 1990;117(3):331-334 Doi: [https://www.doi.org/10.1016/0304-3940\(90\)90686-4](https://www.doi.org/10.1016/0304-3940(90)90686-4)
 19. M. J. Caterina, M. A. Schumacher, M. Tominaga, T. A. Rosen, J. D. Levine and D. Julius. **The capsaicin receptor: a heat-activated ion channel in the pain pathway.** *Nature* 1997;389(6653):816-824 Doi: <https://www.doi.org/10.1038/39807>
 20. M. Tominaga, M. J. Caterina, A. B. Malmberg, T. A. Rosen, H. Gilbert, K. Skinner, B. E. Raumann, A. I. Basbaum and D. Julius. **The cloned capsaicin receptor integrates multiple pain-producing stimuli.** *Neuron* 1998;21(3):531-543 Doi: [https://www.doi.org/10.1016/s0896-6273\(00\)80564-4](https://www.doi.org/10.1016/s0896-6273(00)80564-4)
 21. L. K. Singh, X. Pang, N. Alexacos, R. Letourneau and T. C. Theoharides. **Acute immobilization stress triggers skin mast cell degranulation via corticotropin releasing hormone, neurotensin, and substance P: A link to neurogenic skin disorders.** *Brain Behav Immun* 1999;13(3):225-239 Doi: <https://www.doi.org/10.1006/brbi.1998.0541>
 22. R. L. Baylie and J. E. Brayden. **TRPV channels and vascular function.** *Acta Physiol (Oxf)* 2011;203(1):99-116 Doi: <https://www.doi.org/10.1111/j.1748-1716.2010.02217.x>
 23. J. J. C. s. Szolcsanyi. **Capsaicin, irritation, and desensitization: neurophysiological basis and future perspectives.** 1990;2(141-168
 24. M. Dux, C. Will, B. Vogler, M. R. Filipovic and K. Messlinger. **Meningeal blood flow is controlled by H2 S-NO crosstalk activating a HNO-TRPA1-CGRP signalling pathway.** *Br J Pharmacol* 2016;173(3):431-445 Doi: <https://www.doi.org/10.1111/bph.13164>
 25. M. N. Sullivan, A. L. Gonzales, P. W. Pires, A. Bruhl, M. D. Leo, W. Li, A. Oulidi, F. A. Boop, Y. Feng, J. H. Jaggar, D. G. Welsh and S. Earley. **Localized TRPA1 channel Ca²⁺ signals stimulated by reactive oxygen species promote cerebral artery dilation.** *Sci Signal* 2015;8(358):ra2 Doi: <https://www.doi.org/10.1126/scisignal.2005659>
 26. T. C. Theoharides, C. Spanos, X. Pang, L. Alferes, K. Ligris, R. Letourneau, J. J. Rozniecki, E. Webster and G. P. Chrousos. **Stress-induced intracranial mast cell degranulation: a corticotropin-releasing hormone-mediated effect.** *Endocri-*

- nology* 1995;136(12):5745-5750 Doi: <https://www.doi.org/10.1210/endo.136.12.7588332>
27. P. Huang, P. H. Kuo, M. T. Lee, L. C. Chiou and P. C. Fan. **Age-Dependent Anti-migraine Effects of Valproic Acid and Topiramate in Rats.** *Front Pharmacol* 2018;9(1):1095 Doi: <https://www.doi.org/10.3389/fphar.2018.01095>
 28. V. Dimitriadou, M. G. Buzzi, M. A. Moskowitz and T. C. Theoharides. **Trigeminal sensory fiber stimulation induces morphological changes reflecting secretion in rat dura mater mast cells.** *Neuroscience* 1991;44(1):97-112 Doi: [https://www.doi.org/10.1016/0306-4522\(91\)90253-k](https://www.doi.org/10.1016/0306-4522(91)90253-k)
 29. V. Dimitriadou, A. Rouleau, M. D. Trung Tuong, G. J. Newlands, H. R. Miller, G. Luffau, J. C. Schwartz and M. Garbarg. **Functional relationships between sensory nerve fibers and mast cells of dura mater in normal and inflammatory conditions.** *Neuroscience* 1997;77(3):829-839 Doi: [https://www.doi.org/10.1016/s0306-4522\(96\)00488-5](https://www.doi.org/10.1016/s0306-4522(96)00488-5)
 30. I. Shelukhina, N. Mikhailov, P. Abushik, L. Nurullin, E. E. Nikolsky and R. Giniatullin. **Cholinergic Nociceptive Mechanisms in Rat Meninges and Trigeminal Ganglia: Potential Implications for Migraine Pain.** *Front Neurol* 2017;8(1):163 Doi: <https://www.doi.org/10.3389/fneur.2017.00163>
 31. J. Cheng, X. N. Yang, X. Liu and S. P. Zhang. **Capsaicin for allergic rhinitis in adults.** *Cochrane Database Syst Rev* 2006;2(1):Cd004460 Doi: <https://www.doi.org/10.1002/14651858.CD004460.pub2>
 32. S. Frydas, G. Varvara, G. Murmura, A. Saggini, A. Caraffa, P. Antinolfi, S. Tete, D. Tripodi, F. Conti, E. Cianchetti, E. Toniato, M. Rosati, L. Speranza, A. Pantalone, R. Saggini, L. M. Di Tommaso, T. C. Theoharides, P. Conti and F. Pandolfi. **Impact of capsaicin on mast cell inflammation.** *Int J Immunopathol Pharmacol* 2013;26(3):597-600 Doi: <https://www.doi.org/10.1177/039463201302600303>
 33. D. Levy, V. Kainz, R. Burstein and A. M. Strassman. **Mast cell degranulation distinctly activates trigemino-cervical and lumbosacral pain pathways and elicits widespread tactile pain hypersensitivity.** *Brain Behav Immun* 2012;26(2):311-317 Doi: <https://www.doi.org/10.1016/j.bbi.2011.09.016>
 34. J. Wallengren and R. Håkanson. **Effects of substance P, neurokinin A and calcitonin gene-related peptide in human skin and their involvement in sensory nerve-mediated responses.** *Eur J Pharmacol* 1987;143(2):267-273 Doi: [https://www.doi.org/10.1016/0014-2999\(87\)90542-5](https://www.doi.org/10.1016/0014-2999(87)90542-5)
 35. T. Boes and D. Levy. **Influence of sex, estrous cycle, and estrogen on intracranial dural mast cells.** *Cephalalgia* 2012;32(12):924-931 Doi: <https://www.doi.org/10.1177/0333102412454947>
 36. M. Baun, M. H. Pedersen, J. Olesen and I. Jansen-Olesen. **Dural mast cell degranulation is a putative mechanism for headache induced by PACAP-38.** *Cephalalgia* 2012;32(4):337-345 Doi: <https://www.doi.org/10.1177/0333102412439354>